



# Genomic, proteomic and lipidomic evaluation of endometrial receptivity

## Endometrial reseptivitenin genomik, proteomik ve lipidomik deęerlendirmesi

İrem Demiral<sup>1</sup>, Murat Doęan<sup>2</sup>, Ercan Bastu<sup>1</sup>, Faruk Buyru<sup>1</sup>

<sup>1</sup>Istanbul University Faculty of Medicine, Department of Obstetrics and Gynecology, Istanbul, Turkey

<sup>2</sup>Acibadem Fulya Hospital, Division of Reproductive Endocrinology and Infertility, Istanbul, Turkey

### Abstract

Endometrial receptivity is a complex phenomenon that plays a vital role in infertility. Although quality of embryo can be evaluated for a successful implantation, endometrial receptivity is still an unknown factor. With advances in technology, the microarray approach has provided an 'omic' tool to evaluate endometrial receptivity. In Latin, 'omic' means the whole family. The genomic, proteomic, and lipidomic evaluations of endometrium mean a wholesome evaluation of the genes, lipids and proteins of the endometrium. Evaluation of receptivity with this three-way approach may provide insight to the potential markers of implantation. Genomic analysis has been limited to date because not every gene alteration affects protein expression. Lipidomic analysis has recently gained popularity because lipids are strictly controlled during the implantation period. In summary, with the recent advances in microarray technology, genomic, lipidomic, and proteomic analyses of the endometrium may provide 'optimal' evaluation tools and criteria to assess receptivity in the near future.

**Keywords:** Endometrial receptivity, genomic, lipidomic, proteomic

### Öz

Endometrial reseptivite, bugün hala gizemi net olarak çözülememiş olan ancak tedavi umudu içinde olan infertil çiftlerde aydınlatılmasını önemli, belki de en hayati ehemmiyete sahip olan noktadır. Başarılı bir implantasyon için embriyonun kalitesi deęerlendirilebilir iken, endometrial reseptivite bilinmez kalmaktadır. Her geen gün daha da ilerleyen microarray teknolojisi, endometrial reseptivitenin deęerlendirilmesinde 'omik' bakış açısını getirmiştir. 'Omik', bir ailenin bütünü anlamına gelmektedir. Endometriyumun genomik, proteomik ve lipidomik araştırılması, endometriyumun genlerinin, proteinlerin ve lipidlerinin tümünün alışılmasıdır. Reseptivitenin bu üç başlık altında araştırılması, implantasyona olanak saęlayan markerların belirlenmesinde önemli bir adımdır. Endometriyumda her gen deęişikliği, protein ekspresyonunda deęişikliğine neden olmadığından, genomik araştırmaların daha geri planda olmasına neden olmuştur. İmplantasyon aralığı boyunca, lipidlerin sıkı kontrol altında olmasının anlaşılması, lipidomik araştırmaların üzerinde daha fazla durulmasına neden olmuştur. Özetle, endometrial reseptivitenin microarray teknolojisi ışığında genomik, proteomik ve lipidomik açıdan araştırılması, ilerleyen yıllarda, 'optimal' olarak deęerlendirilebilecek bir endometriyum için gerekli olan kriterlerin belirlenilmesini saęlayacağına inanılmaktadır.

**Anahtar Kelimeler:** Endometrial reseptivite, genomic, lipidomik, proteomik

### Introduction

Three important events have to occur during the window of implantation in order to form a healthy pregnancy:(1) proper embryo development(2) development of a receptive endometrium, and(3) a successful interaction between the embryo and endometrium. Even when optimal conditions were met, the pregnancy rate was found to be around 30%(1).

The endometrium undergoes cyclic morphologic and molecular changes in preparation for pregnancy. Understanding these changes is important in order to clarify the basic biology of the uterus and the environment of the endometrial tissue that

allows implantation. Recent studies showed that the uterine environment allows implantation and provides adequate nutrition to the embryo to ensure its development(2,3).

Despite all the advances in reproductive medicine, in vitro fertilization (IVF) success rates are still low. It is thought that the reason of low pregnancy rates in IVF cycles is largely due to problems in endometrial receptivity. In order to access the endometrial receptivity, many parameters such as endometrial morphology or subendothelial blood flow have been used, but their prognostic values were found to be low(4). For the last 20 years, regardless of the advances in embryo quality

Address for Correspondence/Yazışma Adresi: Ercan Bastu, MD, Istanbul University Faculty of Medicine, Department of Obstetrics and Gynecology, Istanbul, Turkey  
Phone: +90 532 413 41 95 E-mail: ercan.bastu@istanbul.edu.tr

Received /Geliş Tarihi : 29.09.2014

Accepted/Kabul Tarihi : 02.01.2015

improvement and 'best' embryo selection, the progress to a better understanding of endometrial function has lagged behind the biology of the blastocyst. Even though the endometrium can be assessed in a non-invasive way thanks to the current imaging methods, endometrial receptivity is still based upon morphologic features, which has no clinical reflection<sup>(5,6)</sup>. As a result, alternative diagnostic methods to replace or accompany conventional methods are needed. Today, we know that the endometrium changes its normal structure during the IVF cycles as a result of ovarian stimulation<sup>(7,8)</sup>. The endometrium responds to exposure of high progesterone levels with the premature injection of human chorionic gonadotropin, used instead of luteinizing hormone (LH), by changing its constitution. When stimulated cycles were compared with normal cycles, a significant difference in cytokine levels were reported<sup>(9,10)</sup>. This is why markers that reflect endometrial receptivity are needed; patients will not be exposed to expensive IVF treatments and better success rates will be achieved.

With the recently increasing use of microarray technology,<sup>(11)</sup> it is possible to analyze thousands of genes in one particular sample. By courtesy of this technology, we now have the opportunity to analyze the genomics of endometrial development. Additionally, in recent years, with the progression seen in the domain called 'omics,' extensive molecular comparison of cells and tissues has been made possible. With the help of this technologic evolution in reproductive medicine, the opportunity to reveal the molecular signature of many cells and organs related to reproduction has arrived. During the stages of the menstrual cycle, by analyzing different expression of genes in the endometrium with microarray, diagnostic markers that reveal the implantation period can be found. With the aid of proteomic analyzers, proteins that have effects on reproduction can be discovered. In this context, progress by definition has occurred in the 'omics' world: genomics defines the analysis of genes, 'proteomics' defines the analysis of proteins, and also functional classifications (such as secretomics, metabolomics, transcriptomics, epigenomics) have been made. In light of this new classification, secretomics is the global definition of factors secreted by cells or organs in physiologic or pathologic conditions<sup>(12,13)</sup>. If we look at the correlation with the uterus, endometrial fluid or tissue samples can be collected using minimal invasive methods and by doing so, endometrial receptivity can be analyzed.

The aim of this review was to focus on the potential advantages of investigating endometrial receptivity, an area that has been studied the least in reproductive medicine, and to scrutinize the clinical implications for IVF in view of current methods of analysis such as genetic expression (genomics), secretomics, proteomics, and lipidomics.

### **Endometrial environment and reproduction**

In humans, the endometrium undergoes two phases during the menstrual cycle: the proliferative and secretory phases.

In each phase, endometrial tissue has its own morphologic and physiologic features. In the middle of secretory phase, the endometrial secretions that fill the uterine lumen interact with the blastocyst. This is called the window of implantation. Special interest is given to these secretions for two reasons: the first is the molecular content of these secretions may be a marker for the endometrium to allow implantation. This situation could have a very important diagnostic value for IVF. The second revolves around the exact molecular environment when the blastocyst enters the uterine lumen; this would give us information about how the endometrium becomes receptive to implantation and amenable to blastocyst development.

The role of the endometrial fluid was recently studied for its capacity to provide adequate nutrition to the developing embryo. This situation is crucial for implantation as much as for the subsequent embryo development. According to animal studies, changes in the mother's diet can lead to adverse outcomes in infants<sup>(14-17)</sup>. Before implantation, it is understood that the embryo makes changes in order to adapt to the environment, cellular distribution between trophoblast and inner cell mass, function of mitochondria, and cellular energy sensors and signal system sensitivity, each of which is associated with metabolism (amino acid cycle, carbohydrate metabolism, nutrition transport)<sup>(14,18-21)</sup>.

### **Analysis of endometrial receptivity**

Endometrial receptivity is mostly dependent on the local effects of estrogen and progesterone<sup>(22)</sup>. These effects, through stimulation of the cellular and molecular responses of the endometrium, can be defined as the endometrial receptivity window, which occurs between the 19<sup>th</sup> and 23<sup>rd</sup> day of the menstrual cycle. For more than 50 years, the gold standard of clinical assessment of the endometrium has been the histologic criteria described by Noyes et al.<sup>(23)</sup>. These investigators described the specific morphologic appearance of the different parts of the endometrium during the menstrual cycle. The classic work of Noyes et al.<sup>(23)</sup> has been cited thousands of times in the literature and has been accepted worldwide as the diagnostic tool in endometrial research. However, in the last ten years, some randomized studies reported that the histologic analysis per se may not be sufficient to predict receptivity<sup>(5,6)</sup>. The fact underlined by these studies was the need for new technologies that will enable objective analyses of endometrial samples and development.

Among the new technologies, microarray is undoubtedly the most used one in reproductive medicine. Ponnampalam et al.<sup>(24)</sup> were the first investigators to reveal the genomic character of the endometrium during the menstrual cycle using microarray technology. These investigators reported that it was possible to stage the endometrium very acutely according to the transcriptional profile, independent of the morphologic appearance. Most importantly, by identifying characteristics of variant genes in different phases of the menstrual cycle,

they showed that gene expression profiling was a potential instrument in the analysis of endometrial receptivity.

Some studies reported that in cycles where ovarian stimulation was performed, high estrogen and/or progesterone levels deteriorated endometrial receptivity<sup>(25,26)</sup>. High estrogen concentrations may lead to early endometrial secretion by previous stimulation of progesterone receptors in the endometrium. Researchers have reported that GnRH agonists and antagonists have little effect on receptivity compared with natural cycles,<sup>(27,28)</sup> and others reported a profound impact<sup>(29-34)</sup>. Steps that require investigation in the future are the similarities or differences of the effects of GnRH agonists and antagonists on endometrial receptivity.

As an alternative to genomics analysis, proteomic and/or lipidomic analysis of uterine fluid and tissue samples can give precious information about molecular physiology<sup>(35,36)</sup>. During IVF, this kind of small-sample-size and quick-result technology can provide a sensitive and accurate assessment of endometrial receptivity through easy samples collection from the uterus via a minimally invasive approach. Research conducted in the last ten years has shown that fluid aspiration performed before embryo transfer can be performed with no adverse effect on pregnancy rates in IVF cycles<sup>(37,38)</sup>.

### Genomic analysis of the endometrium

With the development of new microarray technologies, many investigators have started genomic analysis in reproductive medicine<sup>(39-43)</sup>. Many important molecules have been identified with these studies: genes responsible for lipid metabolism (apolipoprotein D), immune response (decay accelerating factor-CD55, serine or cysteine proteinase, interleukin (IL)-15), control of the cell cycle (cell growth arrest and TNF- $\alpha$ -induced DNA damage), or enzymes whose function differs in different tissues (monoamine oxidase). There are many studies in the literature about the changes in the endometrium's gene expression profile during the transition from the pre-receptive- (LH+1/5) to the receptive period (LH+7/9)<sup>(39,44-47)</sup>. The number of genes whose expression differs between these two periods varies from 107 to 2878.

These genes, which might be the specific markers of endometrial receptivity, were identified in normal menstrual cycles. It is important to know if these genes have clinical value in predicting pregnancy in stimulated cycles. In a recently published study, the endometrial gene expression profile was analyzed in patients for whom ICSI treatment was planned at LH+7/9, 1-2 cycles before starting GnRH analogs. In patients who became pregnant, only 6 genes were found to have increased expression in the samples: vascular endothelial growth factor A (VEGFA), phospholipase A2 group 2A (PLA2G2A), alkaline phosphatase (ALP), leukemia inhibitory factor (LIF), nicotinamide N-methyltransferase (NNMT) and stanniocalcin 1 (STC1)<sup>(48)</sup>. Extensive cohort studies are needed to identify the predictive value of these genes for pregnancy outcomes.

The changes of gene expression seen in the endometrium does not always result in changes in proteins. The genomic analysis of the endometrium is insufficient to understand and identify receptivity in correlation with secreted proteins and other agents; this led to a focus on proteomic and lipidomic analysis. An interesting result of genomic analysis was the revealing of continuous control on lipid metabolism during endometrial development and the implantation window<sup>(40)</sup>. In this context, the importance of lipidomic analysis of endometrial tissue and fluids is underlined.

### Proteomic analysis of the endometrium

As stated before, there is a significant increase in the protein content of fluids that fill the uterine lumen during the implantation window. As such, the explicit protein profile of uterine fluid in fertile women may play an important role in identifying the receptive endometrium. Proteomic analysis may help to predict the right implantation window or to determine how much of the endometrium is 'open' to receptivity during IVF approaches<sup>(49-54)</sup>. One common protein has been reported in studies where proteomic analysis of the endometrium was performed from tissue samples: annexin A4<sup>(44,46)</sup>. As previously discussed, the genomic changes do not always correlate with proteomic changes. For this reason, the results of six transcriptomic<sup>(39,42-47)</sup> and two proteomic<sup>(44,46)</sup> studies conducted during the transition of the endometrium from the pre-receptive phase to the receptive phase were compared. Only gene of one protein (annexin A4) was found to be up-regulated in five transcriptomic and both proteomic studies<sup>(42-47)</sup>. This protein is thought to play a role in shaping the endometrial apical pole for cellular adhesion. Annexin A2 (ANXA2) and monoamine oxidase A (MAOA) genes were found over-expressed in three transcriptomic studies: ANXA2<sup>(42,43,47)</sup>, MAOA<sup>(42,45,46)</sup> and one proteomic study<sup>(44)</sup>. It is shown in the literature that patients who experience implantation failures have decreased MAOA mRNA and protein expression<sup>(55,56)</sup>. Transgelin 2 (TAGLN, TGF- $\beta$ /Smad-dependent epithelial cell migration) and L-plastin (LCPI, a member of the actine-binding protein family) were found to be over-expressed in two transcriptomic studies: TAGLN<sup>(42,46)</sup>, LCPI<sup>(43,47)</sup> and one proteomic study<sup>(44)</sup>. From all these comparisons, information about five more genes was obtained; progesterone-receptor-membrane component 1 (PGRMC 1) and stathmin 1 (STMN) was found down-regulated in one transcriptomic study and this finding was supported in a proteomic study: PGRMC 1<sup>(44,47)</sup>, STMN<sup>(43,44)</sup>. Apolipoprotein L2 (APOL2), an aldehyde dehydrogenase 1 family member, A3 (ALDH1A3), and S100-calcium-binding protein A10 (S100A10) were found to be up-regulated in a transcriptomic study, and a proteomic study supported this finding<sup>(44,46)</sup>. The function of these five genes is as yet undiscovered.

There are some advantages in performing proteomic analysis in samples taken from endometrial fluid compared with tissue

biopsy: the lower content of cellular protein in endometrial fluid makes the analysis more detailed and facilitates the identification of proteins and low-concentration post-translational modifications. The basic limitation is the lack of consensus on the collection of the fluid, which differs between studies.

The protein content of endometrial fluid was first studied in 1998 by Beier and Beier-Hellwig<sup>(57)</sup>. In 2009, Chen et al.<sup>(58)</sup> reported 196 different proteins secreted from the endometrium during the mid-proliferative phase compared with the mid-secretory phase.

Proteomic analysis of uterine secretions has been the subject of much research<sup>(49,52,54)</sup>. In summary, the information gathered from these studies shows us that the main functions of the proteins found in uterine secretions are especially related with cytoskeleton regulation, cellular adhesion, protein folding, and signal transduction.

### Lipidomic analysis of the endometrium

Compared with genomic and proteomic analysis, lipidomic analysis has thus far been the least used method among all the research on the endometrium<sup>(59)</sup>. On the other hand, we know that the endometrium has a very important lipid component for reproduction. Triglycerides and eicosanoids can be counted among the lipid mediators secreted from the endometrium. From the eicosanoid family, prostaglandins, thromboxanes, leukotrienes, endocannabinoids and sphingolipids have been observed to play a role in reproduction<sup>(60-64)</sup>. Among them, endocannabinoids especially, lysophosphatidic acid (LPA) and prostaglandins (PG) were the most studied agents<sup>(65)</sup>. The consistency of results of research conducted on different animal models supports the hypothesis that lipids could be critical for implantation.

Two basic endocannabinoids were found to play a role in implantation in rodents: anandamide N-arachidonoyl ethanolamine (AEA) and 2-arachidonoylglycerol. Abnormal levels of these two lipids lead to the postponing of implantation and a decrease in pregnancy outcomes<sup>(66)</sup>. The basic enzyme that degrades endocannabinoids is fatty acid amide hydrolase (FAAH) and it was understood that this enzyme was associated with high pregnancy rates after IVF and embryo transfer (ET)<sup>(65)</sup>. It is reported that low FAAH levels leads to an increase in AEA and subsequent decrease in uterine receptivity<sup>(67)</sup>.

Lysophosphatidic acid is a water-soluble phospholipid. It is a signal transduction molecule that has many functions in different organs. In a study conducted in mice, LPA was shown to be necessary for normal embryo size by stimulating its receptor LPA3,<sup>(61)</sup> which is a positive factor for implantation. According to a recent study, lysophosphatidic acid regulates the endometrium for implantation during the receptive period by controlling levels of endocannabinoid and prostaglandin mediators via its receptor LPA3<sup>(64,65)</sup>.

Prostaglandins (PG) are the final products of arachidonic acid metabolism. Phospholipids present in membranes turn into PG

with phospholipase A2 (PLA2) and COX-2. With these enzymes, PG levels in the lumen and stroma increase. Animal studies reported interesting results about the effects of PG and COX enzymes on reproduction. For example, it was observed in mice that the PG increase is limited in the area where the blastocyst implants<sup>(68)</sup>. In that study, pregnancy in mice was totally 'prevented' with the inhibition of COX-1 and COX-2, thus showing the importance of prostaglandins during pregnancy<sup>(68)</sup>. The prevention of pregnancy was restored partially with the exogenous application of PGI2 and PGE2. In this case, prostaglandins must play an important role in embryo implantation in mice. In another study, PGE2 was found increased in the implantation area, whereas PGF2- $\alpha$  was mostly found between implantation areas<sup>(69)</sup>. However, in 2006, Cong et al.<sup>(70)</sup> reported that when the implantation period was compared with other periods of the menstrual cycle, PGF2- $\alpha$  was not increased; the highest increase was seen in PGI2 followed by PGE2. PGI2 and PGE2 is known to increase the vascular permeability, which is necessary for ovulation and the start of implantation<sup>(71)</sup>. However, in one study conducted on mice, the mediator that increased vascular permeability in the implantation area was found to be PGE2, not PGI2<sup>(59)</sup>. PGE2 was found to play a positive role in implantation in different kinds animals<sup>(60,72,73)</sup>. In 2013, Vilella et al.<sup>(74)</sup> collected 51 endometrial fluid samples from women and observed that PGE2 and PGF2 $\alpha$  levels were significantly increased during the implantation window.

In a recent study, a relation between defective endometrial prostaglandin synthesis and repeated implantation failure was established in women undergoing IVF treatment<sup>(75)</sup>. This study showed that prostaglandins were important mediators at early-stage pregnancy.

### Conclusion

Endometrial receptivity is a crucial process during embryo implantation. The common result of all the genomic analyses mentioned in this review is that endometrial receptivity is a complex process that involves countless genes. Some studies report that some genes are continuously expressed in the endometrium and modifications in these genes can help us to better understand endometrial receptivity.

Proteomic and lipidomic research has started to find its place in reproductive medicine. It will take time for the results of research to become a specialized tool for diagnosis and treatment. At the same time, even potential suppositions are exciting. For example, drugs that women use during IVF treatment might change the molecular composition of endometrial tissue and secretions by altering the physiology of the endometrium. With the help of proteomic and lipidomic analysis, objective treatment protocols developed according to the patient's personal features can be identified. In this context, genomic, proteomic, and lipidomic analysis will probably have to be used together.

In conclusion, the genomic, proteomic, and lipidomic analysis of endometrial tissues and secretions can provide investigators with important information about how genes, protein, and

lipid mediators work and how they can enable endometrial receptivity ready for embryo implantation. Thus, effective treatment protocols can be developed in order to increase the basic aim of IVF strategies, the pregnancy rate. The subsequent step is testing the endometrium in terms of receptivity and guiding modifications of ovarian stimulation and luteal phase support protocols.

### Authorship Contributions

*Concept:* Ercan Baştu, *Design:* Ercan Baştu, *Data Collection:* İrem Demiral, *Analysis:* İrem Demiral, *Literature Search:* Murat Doğan, *İrem Demiral*, *Writing:* Murat Doğan, *İrem Demiral*, *Peer-review:* Externally peer-reviewed, *Conflict of Interest:* No conflict of Interest was declared by authors, *Financial Disclosure:* The authors declared that this study has received no financial support.

### References

- Macklon NS, Geraedts JP, Fauser BC. Conception to ongoing pregnancy: the 'black box' of early pregnancy loss. *Hum Reprod Update* 2002;8:333-43.
- Fleming TP, Kwong WY, Porter R, Ursell E, Fesenko I, Wilkins A, et al. The embryo and its future. *Biol Reprod* 2004;71:1046-54.
- Fleming TP, Velazquez MA, Eckert JJ, Lucas ES, Watkins AJ. Nutrition of females during the peri-conceptual period and effects on foetal programming and health of offspring. *Anim Reprod Sci* 2012;130:193-7.
- Aghajanova L, Hamilton AE, Giudice LC. Uterine receptivity to human embryonic implantation: histology, biomarkers biomarkers, and transcriptomics. *Semin Cell Dev Biol* 2008;19:204-11.
- Coutifaris C, Myers ER, Guzick DS, Diamond MP, Carson SA, Legro RS, et al. Histological dating of timed endometrial biopsy tissue is not related to fertility status. *Fertil Steril* 2004;82:1264-72.
- Murray MJ, Meyer WR, Zaino RJ, Lessey BA, Novotny DB, Ireland K, et al. A critical analysis of the accuracy, reproducibility, and clinical utility of histologic endometrial dating in fertile women. *Fertil Steril* 2004;81:1333-43.
- Devroey P, Bourgain C, Macklon NS, Fauser BC. Reproductive biology and IVF: ovarian stimulation and endometrial receptivity. *Trends Endocrinol Metab* 2004;15:84-90.
- Kolibianakis EM, Bourgain C, Platteau P, Albano C, Van Steirteghem AC, Devroey P. Abnormal endometrial development occurs during the luteal phase of nonsupplemented donor cycles treated with recombinant follicle-stimulating hormone and gonadotropin-releasing hormone antagonists. *Fertil Steril* 2003;80:464-6.
- Haouzi D, Dechaud H, Assou S, De Vos J, Hamamah S. Insights into human endometrial receptivity from transcriptomic and proteomic data. *Reprod Biomed Online* 2012;24:23-34.
- Horcajadas JA, Sharkey AM, Catalano RD, Sherwin JRA, Dominguez F, Burgos LA, et al. Effect of an intrauterine device on the gene expression profile of the endometrium. *J Clin Endocrinol Metab* 2006;91:3199-207.
- Schena M, Shalon D, Davis RW, Brown PO. Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 1995;270:467-70.
- Dominguez F, Gadea B, Mercader A, Esteban FJ, Pellicer A, Simon C. Embryologic outcome and secretome profile of implanted blastocysts obtained after coculture in human endometrial epithelial cells versus the sequential system. *Fertil Steril* 2010;93:774-82.
- Garrido-Gomez T, Dominguez F, Lopez JA, Camafeita E, Quinonero A, Martinez- Conejero JA, et al. Modeling human endometrial decidualization from the interaction between proteome and secretome. *J Clin Endocrinol Metab* 2011;96:706-16.
- Kwong WY, Wild AE, Roberts P, Willis AC, Fleming TP. Maternal undernutrition during the preimplantation period of rat development causes blastocyst abnormalities and programming of postnatal hypertension. *Development* 2000;127:4195-202.
- Kwong WY, Osmond C, Fleming TP. Support for Barker hypothesis upheld in rat model of maternal undernutrition during the preimplantation period: application of integrated 'random effects' statistical model. *Reprod Biomed Online* 2004;8:574-6.
- Kwong WY, Miller DJ, Wilkins AP, Dear MS, Wright JN, Osmond C, et al. Maternal low protein diet restricted to the preimplantation period induces a gender-specific change on hepatic gene expression in rat fetuses. *Mol Reprod Dev* 2007;74:48-56.
- Kwong WY, Miller DJ, Ursell E, Wild AE, Wilkins AP, Osmond C, et al. Imprinted gene expression in the rat embryo-fetal axis is altered in response to periconceptual maternal low protein diet. *Reproduction* 2006;132:265-77.
- Mitchell M, Schulz SL, Armstrong DT, Lane M. Metabolic and mitochondrial dysfunction in early mouse embryos following maternal dietary protein intervention. *Biol Reprod* 2009;80:622-30.
- Eng GS, Sheridan RA, Wyman A, Chi MM, Bibee KP, Jungheim ES, et al. AMP kinase activation increases glucose uptake, decreases apoptosis, and improves pregnancy outcome in embryos exposed to high IGF-I concentrations. *Diabetes* 2007;56:2228-34.
- Wyman A, Pinto AB, Sheridan R, Moley KH. One-cell zygote transfer from diabetic to nondiabetic mouse results in congenital malformations and growth retardation in offspring. *Endocrinology* 2008;149:466-9.
- Van Winkle LJ, Tesch JK, Shah A, Campione AL. System B0,+ amino acid transport regulates the penetration stage of blastocyst implantation with possible long-term developmental consequences through adulthood. *Hum Reprod Update* 2006;12:145-57.
- Critchley HO, Saunders PT. Hormone receptor dynamics in a receptive human endometrium. *Reprod Sci* 2009;16:191-9.
- Noyes RW, Hertig AT, Rock J. Dating the endometrial biopsy. *Am J Obstet Gynecol* 1975;122:262-3.
- Ponnampalam AP, Weston GC, Trajstman AC, Susil B, Rogers PA. Molecular classification of human endometrial cycle stages by transcriptional profiling. *Mol Hum Reprod* 2004;10:879-93.
- Krikun G, Schatz F, Taylor R, Critchley HO, Rogers PA, Huang J, et al. Endometrial endothelial cell steroid receptor expression and steroid effects on gene expression. *J Clin Endocrinol Metab* 2005;90:1812-8.
- Liu Y, Lee KF, Ng EH, Yeung WS, Ho PC. Gene expression profiling of human peri-implantation endometria between natural and stimulated cycles. *Fertil Steril* 2008;90:2152-64.
- Mirkin S1, Nikas G, Hsiu JG, Diaz J, Oehninger S. Gene expression profiles and structural/functional features of the peri-implantation endometrium in natural and gonadotropin- stimulated cycles. *J Clin Endocrinol Metab* 2004;89:5742-52.
- Simon C, Oberye J, Bellver J, Vidal C, Bosch E, Horcajadas JA, et al. Similar endometrial development in oocyte donors treated with either high- or standard-dose GnRH antagonist compared to treatment with a GnRH agonist or in natural cycles. *Hum Reprod* 2005;20:3318-27.
- Haouzi D, Assou S, Dechanet C, Anahory T, Dechaud H, De Vos J, et al. Controlled ovarian hyperstimulation for in vitro fertilization alters endometrial receptivity in humans: protocol effects. *Biol Reprod* 2010;82:679-86.

30. Horcajadas JA, Riesewijk A, Polman J, van Os R, Pellicer A, Mosselman S, et al. Effect of controlled ovaria hyperstimulation in IVF on endometrial gene expression profiles. *Mol Hum Reprod* 2005;11:195-205.
31. Horcajadas JA, Mínguez P, Dopazo J, Esteban FJ, Dominguez F, Giudice LC, et al. Controlled ovarian stimulation induces a functional genomic delay of the endometrium with potential clinical implications. *J Clin Endocrinol Metab* 2008;93:4500-10.
32. Macklon NS, van der Gaast MH, Hamilton A, Fauser BC, Giudice LC. The impact of ovarian stimulation with recombinant FSH in combination with GnRH antagonist on the endometrial transcriptome in the window of implantation. *Reprod Sci* 2008;15:357-65.
33. Ruan HC, Zhu XM, Luo Q, Liu AX, Qian YL, Zhou CY, et al. Ovarian stimulation with GnRH agonist, but not GnRH antagonist, partially restores the expression of endometrial integrin beta3 and leukaemia-inhibitory factor and improves uterine receptivity in mice. *Hum Reprod* 2006;21:2521-9.
34. Van Vaerenbergh I, Van Lommel L, Ghislain V, In't Veld P, Schuit F, Fatemi HM, et al. In GnRH antagonist/rec-FSH stimulated cycles, advanced endometrial maturation on the day of oocyte retrieval correlates with altered gene expression. *Hum Reprod* 2009;24:1085-91.
35. Urzua MA, Stambaugh R, Flickinger G, Mastroianni Jr L. Uterine and oviduct fluid protein patterns in the rabbit before and after ovulation. *Fertil Steril* 1970;21:860-5.
36. Ulbrich SE, Schulke K, Groebner AE, Reichenbach HD, Angioni C, Geisslinger G, et al. Quantitative characterization of prostaglandins in the uterus of early pregnant cattle. *Reproduction* 2009;138:371-82.
37. Boomsma CM, Kavelaars A, Eijkemans MJ, Lentjes EG, Fauser BC, Heijnen CJ, et al. Endometrial secretion analysis identifies a cytokine profile predictive of pregnancy in IVF. *Hum Reprod* 2009;24:1427-35.
38. van der Gaast MH, Beier-Hellwig K, Fauser BC, Beier HM, Macklon NS. Endometrial secretion aspiration prior to embryo transfer does not reduce implantation rates. *Reprod Biomed Online* 2003;7:105-9.
39. Carson DD, Lagow E, Thathiah A, Al-Shami R, Farach-Carson MC, Vernon M, et al. Changes in gene expression during the early to mid-luteal (receptive phase) transition in human endometrium detected by high-density microarray screening. *Mol Hum Reprod* 2002;8:871-9.
40. Kao LC, Tulac S, Lobo S, Imani B, Yang JP, Germeyer A, et al. Global gene profiling in human endometrium during the window of implantation. *Endocrinology* 2002;143:2119-38.
41. Borthwick JM, Charnock-Jones DS, Tom BD, Hull ML, Teirney R, Phillips SC, et al. Determination of the transcript profile of human endometrium. *Mol Hum Reprod* 2003;9:19-33.
42. Riesewijk A, Martin J, van Os R, Horcajadas JA, Polman J, Pellicer A, et al. Gene expression profiling of human endometrial receptivity on days LH+2 versus LH+7 by microarray technology. *Mol Hum Reprod* 2003;9:253-64.
43. Mirkin S, Arslan M, Churikov D, Corica A, Diaz JI, Williams S, et al. In search of candidate genes critically expressed in the human endometrium during the window of implantation. *Hum Reprod* 2005;20:2104-17.
44. Domínguez F, Garrido-Gómez T, López JA, Camafeite E, Quiñonero A, Pellicer A, Simón C. Proteomic analysis of the human receptive versus non-receptive endometrium using differential in-gel electrophoresis and MALDI-MS unveils stathmin 1 and annexin A2 as differentially regulated. *Hum Reprod* 2009;24:2607-17.
45. Díaz-Gimeno P, Horcajadas JA, Martínez-Conejero JA, Esteban FJ, Alama P, Pellicer A, et al. A genomic diagnostic tool for human endometrial receptivity based on the transcriptomic signature. *Fertil Steril* 2011;95:50-60, 60.e1-15.
46. Haouzi D, Assou S, Mahmoud K, Tondeur S, Rème T, Hedon B, et al. Gene expression profile of human endometrial receptivity: comparison between natural and stimulated cycles for the same patients. *Hum Reprod* 2009;24:1436-45.
47. Talbi S, Hamilton AE, Vo KC, Tulac S, Overgaard MT, Dosiou C, et al. Molecular phenotyping of human endometrium distinguishes menstrual cycle phases and underlying biological processes in normo-ovulatory women. *Endocrinology* 2006;147:1097-121.
48. Allegra A, Marino A, Coffaro F, Lama A, Rizza G, Scaglione P, et al. Is there a uniform basal endometrial gene expression profile during the implantation window in women who became pregnant in a subsequent ICSI cycle? *Hum Reprod* 2009;24:2549-57.
49. Scotchie JG, Fritz MA, Mocanu M, Lessey BA, Young SL. Proteomic analysis of the luteal endometrial secretome. *Reprod Sci* 2009;16:883-93.
50. Aghajanova L. Leukemia inhibitory factor and human embryo implantation. *Ann N Y Acad Sci* 2004;1034:176-83.
51. Ametzazurra A, Matorras R, Garcia-Velasco JA, Prieto B, Simon L, Martinez A, et al. Endometrial fluid is a specific and non-invasive biological sample for protein biomarker identification in endometriosis. *Hum Reprod* 2009;24:954-65.
52. Casado-Vela J, Rodriguez-Suarez E, Iloro I, Ametzazurra A, Alkorta N, Garcia-Velasco JA, et al. Comprehensive proteomic analysis of human endometrial fluid aspirate. *J Proteome Res* 2009;8:4622-32.
53. Olivennes F, Ledee-Bataille N, Samama M, Kadoch J, Taupin JL, Dubanchet S, et al. Assessment of leukemia inhibitory factor levels by uterine flushing at the time of egg retrieval does not adversely affect pregnancy rates with in vitro fertilization. *Fertil Steril* 2003;79:900-4.
54. Hannan NJ, Stephens AN, Rainczuk A, Hincks C, Rombauts LJ, Salamonsen LA. 2D-DiGE analysis of the human endometrial secretome reveals differences between receptive and nonreceptive states in fertile and infertile women. *J Proteome Res* 2010;9:6256-64.
55. Henriquez S, Tapia A, Quezada M, Vargas M, Cardenas H, Rios M, et al. Deficient expression of monoamine oxidase A in the endometrium is associated with implantation failure in women participating as recipients in oocyte donation. *Mol Hum Reprod* 2006;12:749-54.
56. Tapia A, Gangi LM, Zegers-Hochschild F, Balmaceda J, Pommer R, Trejo L, et al. Differences in the endometrial transcript profile during the receptive period between women who were refractory to implantation and those who achieved pregnancy. *Hum Reprod* 2008;23:340-51.
57. Beier HM, Beier-Hellwig K. Molecular and cellular aspects of endometrial receptivity. *Hum Reprod Update* 1998;4:448-58.
58. Chen JI, Hannan NJ, Mak Y, Nicholls PK, Zhang J, Rainczuk A, et al. Proteomic characterization of midproliferative and midsecretory human endometrium. *J Proteome Res* 2009;8:2032-44.
59. Wenk MR. The emerging field of lipidomics. *Nat Rev Drug Discov* 2005;4:594-610.
60. Wang H, Dey SK. Lipid signaling in embryo implantation. *Prostaglandins Other Lipid Mediat* 2005;77:84-102.
61. Ye X, Hama K, Contos JJ, Anliker B, Inoue A, Skinner MK, et al. LPA3-mediated lysophosphatidic acid signalling in embryo implantation and spacing. *Nature* 2005;435:104-8.
62. Maccarrone M. Endocannabinoids: friends and foes of reproduction. *Prog Lipid Res* 2009;48:344-54.

63. Paria BC, Ma W, Tan J, Raja S, Das SK, Dey SK, et al. Cellular and molecular responses of the uterus to embryo implantation can be elicited by locally applied growth factors. *Proc Natl Acad Sci U S A* 2001;98:1047-52.
64. Mizugishi K, Li C, Olivera A, Bielawski J, Bielawska A, Deng CX, et al. Maternal disturbance in activated sphingolipid metabolism causes pregnancy loss in mice. *J Clin Invest* 2007;117:2993-3006.
65. Sordelli MS, Beltrame JS, Cella M, Gervasi MG, Perez Martinez S, Burdet J, et al. Interaction between lysophosphatidic acid, prostaglandins and the endocannabinoid system during the window of implantation in the rat uterus. *PLoS One* 2012;7:e46059.
66. Maccarrone M, DeFelici M, Klinger FG, Battista N, Fezza F, Dainese E, et al. Mouse blastocysts release a lipid which activates anandamide hydrolase in intact uterus. *Mol Hum Reprod* 2004;10:215-21.
67. Schmid PC, Paria BC, Krebsbach RJ, Schmid HH, Dey SK. Changes in anandamide levels in mouse uterus are associated with uterine receptivity for embryo implantation. *Proc Natl Acad Sci U S A* 1997;94:4188-92.
68. Reese J, Zhao X, Ma WG, Brown N, Maziasz TJ, Dey SK. Comparative analysis of pharmacologic and/or genetic disruption of cyclooxygenase-1 and cyclooxygenase-2 function in female reproduction in mice. *Endocrinology* 2001;142:3198-206.
69. Pakrasi PL. Prostaglandins and ovum implantation in mice. *J Exp Zool* 1997;278:53-7.
70. Cong J, Diao H-L, Zhao Y-C, Ni H, Yan Y-Q, Yang Z-M. Differential expression and regulation of cyclooxygenases, prostaglandin E synthases and prostacyclin synthase in rat uterus during the peri-implantation period. *Reproduction* 2006;131:139-51.
71. Lim H, Paria BC, Das SK, Dinchuk JE, Langenbach R, Trzaskos JM, et al. Multiple female reproductive failures in cyclooxygenase 2-deficient mice. *Cell* 1997;91:197-208.
72. Kennedy TG, Gillio-Meina C, Phang SH. Prostaglandins and the initiation of blastocyst implantation and decidualization. *Reproduction* 2007;134:635-43.
73. Waclawik A, Jabbour HN, Blitek A, Ziecik AJ. Estradiol-17, prostaglandin E2 (PGE2), and the PGE2 receptor are involved in PGE2 positive feedback loop in the porcine endometrium. *Endocrinology* 2009;150:3823-32.
74. Vilella F, Ramirez LB, Simón C. Lipidomics as an emerging tool to predict endometrial receptivity. *Fertil Steril* 2013;99:1100-6.
75. Achache H, Tsafirir A, Prus D, Reich R, Revel A. Defective endometrial prostaglandin synthesis identified in patients with repeated implantation failure undergoing in vitro fertilization. *Fertil Steril* 2010;94:1271-8.